

#### **Lumiprobe Corporation**

201 International Circle, Suite 135 Hunt Valley, Maryland 21030

**USA** 


Phone: +1 888 973 6353 Fax: +1 888 973 6354 Email: order@lumiprobe.com


# FAM phosphoramidite, 5-isomer

http://www.lumiprobe.com/p/5-fam-phosphoramidite

Standard fluorescein (FAM) phosphoramidite for 5'-terminal oligonucleotide labeling, high isomeric purity single isomer. This highly purified amidite reagent ensures excellent coupling results with various synthesizers.

Samples and special offers are available for oligo manufacturers. Contact us for bulk supplies of this reagent.





Structure of FAM Phosphoramidite, 5-Isomer

FAM absorbance and emission spectra

### **General properties**

Appearance: off white solid

Molecular weight: 843.94 CAS number: 147566-42-5 Molecular formula:  $C_{46}H_{58}N_3O_{10}P$ 

Solubility: Good solubility in acetonitrile and DCM

Quality control: NMR  $^{1}$ H and  $^{31}$ P, HPLC-MS (95+%), isomeric purity > 97%

Storage conditions: Storage: 12 months after receival at -20°C in the dark. Transportation: at room

temperature for up to 3 weeks. Avoid prolonged exposure to light. Desiccate.

Legal statement: This Product is offered and sold for research purposes only. It has not been tested for

safety and efficacy in food, drug, medical device, cosmetic, commercial or any other use. Supply does not express or imply authorization to use for any other purpose, including, without limitation, in vitro diagnostic purposes, in the manufacture of food

or pharmaceutical products, in medical devices or in cosmetic products.

#### Spectral properties

Excitation/absorption maximum, nm: 492  $\epsilon$ , L·mol<sup>-1</sup>·cm<sup>-1</sup>: 72000 Emission maximum, nm: 517 Fluorescence quantum yield: 0.93  $CF_{260}$ : 0.22  $CF_{280}$ : 0.17

## Oligo synthesis details

Diluent: acetonitrile

Coupling conditions: coupling time 10 min

Cleavage conditions: ammonia, 2 h at room temperature

Deprotection conditions:

identical to protected nucleobases; when AMA is used, deblock with ammonia alone for 30 min, then add methylamine  $\,$