



Cyanine dyes

Cyanine dyes are molecules containing polymethine bridge between two nitrogen atoms with a delocalized charge:



Due to their structure, cyanines have outstandingly high extinction coefficients often exceeding $100,000 \text{ Lmol}^{-1}\text{cm}^{-1}$. Different substituents allow to control properties of the chromophore, such as absorbance wavelength, photostability, and fluorescence. For example, absorbance and fluorescence wavelength can be controlled by a choice of polymethine bridge length: longer cyanines possess higher absorbance and emission wavelengths up to near infrared region.


A number of cyanine dyes have been used for life science applications. A series of thiazole and oxazole dyes have been used as DNA- and protein-binding dyes (like TOTO, YOYO, Stains All and others). But the most popular cyanine dyes for life science research were introduced by Alan Waggoner and colleagues of Carnegie Mellon University in early 1990s. The dyes were a modification of cyanine dye Indocyanine Green (ICG) which was used for angiography since 1970s, and they all contain two indolenine rings flanking polymethyne chain. The dyes were found to exhibit low non-specific binding to biomolecules, and have bright fluorescence owing to their huge extinction coefficients and good quantum yields. Once patented, these molecules are now in public domain after expiration of patents of CMU, and are available for purchase from Lumiprobe for research and commercial use as various reactive derivatives, such as [NHS esters](#), [maleimides](#), [azides](#) for Click chemistry, and other derivatives.

There are two varieties of cyanine dyes: non-sulfonated cyanines, and sulfonated cyanines. For many applications they are interchangeable, because their spectral properties are nearly identical. Both sulfonated and non-sulfonated dyes can be used for the labeling of biomolecules such as DNA and proteins. The difference between the dyes is their solubility: sulfo- dyes are water-soluble, and they do not use of organic co-solvent for the labeling in aqueous environment. They are less prone to aggregation in water. There are cases when one of the type of cyanines is desired (see [Sulfonated vs non-sulfonated cyanines](#) section below).

Non-sulfonated cyanines

Available non-sulfonated dyes include Cy3, Cy3.5, Cy5, Cy5.5, Cy7, and Cy7.5. Cy[®] stands for 'cyanine', and the first digit identifies the number of carbon atoms between the indolenine groups. Cy2 which is an oxazole derivative rather than indolenin, is an exception from this rule. The suffix .5 is added for benzo-fused cyanines. Variation of the structures allows to change fluorescence properties of the molecules, and to cover most important part of visible and NIR spectrum with several


fluorophores.



Most derivatives of non-sulfonated cyanines (except for hydrochlorides of hydrazides and amines) have low aqueous solubility. When these molecules are used for biomolecule labeling, use of organic co-solvent (5-20% of DMF or DMSO) is necessary for efficient reaction. Cyanine dye should be dissolved in organic solvent first, and added to a solution of biomolecule (protein, peptide, amino-labeled DNA) in appropriate aqueous buffer. When conjugation takes place efficiently, the dye reacts before it precipitates.

Fluorescent properties of non-sulfonated cyanines have little dependence on solvent and surrounding. Absorbance and fluorescence spectra of non-sulfonated cyanine dyes are plotted below.

ABSORBANCE AND EMISSION SPECTRA OF NON-SULFONATED CYANINE DYES



Lumiprobe Corporation

201 International Circle, Suite 135
Hunt Valley, Maryland 21030
USA
Phone: +1 888 973 6353
Fax: +1 888 973 6354
Email: order@lumiprobe.com

Lumiprobe GmbH

Feodor-Lynen-Strasse 23
30625 Hannover
Germany
Phone: +49 511 16596811
Fax: +49 511 16596815
Email: de@lumiprobe.com


Lumiprobe RUS Ltd

Kotsyubinsky street, 4
121351 Moscow
Russian Federation
Phone: +7 800 775 3271
Email: ru@lumiprobe.com

Sulfonated cyanines

Sulfonated cyanines include additional sulfo-groups which facilitate dissolution of dye molecules in aqueous phase. Charged sulfonate groups decrease aggregation of dye molecules and heavily labeled conjugates.

Currently available sulfonated cyanines include sulfo-Cy3, sulfo-Cy5, and sulfo-Cy7.




Sulfonated cyanines are highly water soluble. No organic co-solvent is needed to perform labeling with these reagents.

Sulfonated vs non-sulfonated cyanines

Sulfonated and non-sulfonated cyanines exhibit very similar fluorescent properties. However, there are a few differences in labeling protocols that should be noticed. Non-sulfonated cyanines must be dissolved in an organic co-solvent (DMF or DMSO) prior to use, and added to a solution of the target molecule in aqueous buffers. The recommended volume of co-solvent should be 10% for Cy3, Cy5, Cy7, and 15% for .5 counterparts. Sulfo-Cyanine reagents can be used in purely aqueous conditions. There is also a difference in purification: when dialysis against water or aqueous buffer is used for purification, sulfo-Cyanine must be used to achieve efficient removal of unreacted dye material. Reactions with both sulfo- and non-sulfo cyanines can be purified by gel filtration, chromatography (HPLC, FPLC, ion exchange), or electrophoresis.

ABSORBANCE AND EMISSION SPECTRA OF SULFONATED VS NON-SULFONATED CYANINES



Sulfonated and non-sulfonated cyanines are interchangeable for the labeling of many classes of targets including:

- soluble proteins, which are tolerant to addition of organic co-solvent

Lumiprobe Corporation

201 International Circle, Suite 135
Hunt Valley, Maryland 21030
USA
Phone: +1 888 973 6353
Fax: +1 888 973 6354
Email: order@lumiprobe.com

Lumiprobe GmbH

Feodor-Lynen-Strasse 23
30625 Hannover
Germany
Phone: +49 511 16596811
Fax: +49 511 16596815
Email: de@lumiprobe.com

Lumiprobe RUS Ltd

Kotsyubinsky street, 4
121351 Moscow
Russian Federation
Phone: +7 800 775 3271
Email: ru@lumiprobe.com

- antibodies (use 5-10% of DMSO/DMF)
- DNA and oligonucleotides
- peptides
- many small molecules

Conjugates produced with similar sulfo- and non-sulfonated reagents (for example, sulfo-Cy5 and Cy5) are very similar in their fluorescent properties, and can be used with various fluorescence instrumentation.

Sulfonated cyanines must be used for:

- sensitive proteins which are denatured by DMF or DMSO
- protein conjugation when purification is done by dialysis
- nanoparticles in aqueous solutions
- insoluble or hydrophobic proteins

Non-sulfonated cyanines must be used for:

- reactions in organic media (dichloromethane, acetonitrile)

The following cyanine products are available from Lumiprobe.

Fluorophore	Reactive form
sulfo-Cy3	NHS ester maleimide azide alkyne carboxylic acid
sulfo-Cy5	NHS ester maleimide azide alkyne carboxylic acid
sulfo-Cy7	NHS ester azide amine carboxylic acid

Cy3	NHS ester maleimide azide alkyne hydrazide amine carboxylic acid
Cy3.5	NHS ester azide carboxylic acid
Cy5	NHS ester maleimide azide alkyne hydrazide amine carboxylic acid
Cy5.5	NHS ester maleimide azide alkyne hydrazide amine carboxylic acid
Cy7	NHS ester maleimide azide alkyne hydrazide amine carboxylic acid
Cy7.5	NHS ester maleimide azide alkyne hydrazide amine carboxylic acid

Cy[®] is a trademark of GE Healthcare.

Lumiprobe Corporation

201 International Circle, Suite 135
 Hunt Valley, Maryland 21030
 USA
 Phone: +1 888 973 6353
 Fax: +1 888 973 6354
 Email: order@lumiprobe.com

Lumiprobe GmbH

Feodor-Lynen-Strasse 23
 30625 Hannover
 Germany
 Phone: +49 511 16596811
 Fax: +49 511 16596815
 Email: de@lumiprobe.com

Lumiprobe RUS Ltd

Kotsyubinsky street, 4
 121351 Moscow
 Russian Federation
 Phone: +7 800 775 3271
 Email: ru@lumiprobe.com