

R6G phosphoramidite, 6-isomer

<http://www.lumiprobe.com/p/6-r6g-phosphoramidite>

R6G phosphoramidite for oligonucleotide synthesis, pure 6-isomer.

R6G (rhodamine 6G) is a xanthene dye of the rhodamine family with high fluorescence quantum yield and high molar extinction coefficient. Relative to those of fluorescein, the absorption (518 nm) and emission (542 nm) maxima of 6-R6G are shifted into the long-wave region.


R6G can be used as a FRET acceptor in DNA-sequencing and in short tandem repeat (STR) analysis. The dye is commonly used for producing TaqMan probes for quantitative PCR. In comparison with HEX and JOE dyes, R6G in combination with BHQ1 quencher has stronger quenching in molecular beacon probes, significantly reducing background fluorescence in PCR.

Usage:


Coupling: 10 minutes.

Deprotection: solution of tert-Butylamine in water (1:3 v/v) overnight at 55°C.

Aqueous ammonia and AMA (solution of 30% ammonium hydroxide/40% aqueous methylamine 1:1 v/v) must NOT be used for deprotection of the modified oligonucleotide from the solid-phase support because of complete and irreversible R6G degradation.



R6G Phosphoramidite structure



General properties

Appearance: Yellowish foam

Molecular weight: 949.93

CAS number: 1355330-47-0

Molecular formula: $C_{46}H_{54}F_6N_5O_8P$

Solubility: good in DCM, acetonitrile

Quality control: NMR 1H and ^{31}P , HPLC-MS (95+%), isomeric purity > 97%

Storage conditions: Storage: 12 months after receipt at -20°C in the dark. Transportation: at room temperature for up to 2 weeks. Avoid prolonged exposure to light. Desiccate.

TN VED Code: 3204190000