Lumiprobe Corporation
115 Airport Dr Suite 160
Westminster, Maryland 21157
USA
Phone: +1 888 973 6353
Fax: +1 888 973 6354
Email: order@lumiprobe.com

Alkyne phosphoramidite, 5'-terminal

http://www.lumiprobe.com/p/alkyne-phosphoramidite-ach

Phosphoramidite for the synthesis of oligonucleotides with 5'-terminal alkyne for click chemistry.

This alkyne amidite has several advantages over 5'-hexynyl phosphoramidite, 5'-butynyl-CEP, and other 5'-terminal alkyne phosphoramidites. First, it is solid compound which is easier to handle and dispense. And due to its structure, it is also more stable in solution, and has longer shelf life

Diluent for this phosphoramidite is acetonitrile, 5 min coupling time is recommended. Because this amidite does not contain 5'-terminal DMT group, no 5'-deprotection needed. Oligonucleotides should be deblocked under standard conditions, and purified by PAGE, or ion exchange HPLC.

Oligonucleotides with this modification are ideal for the use in click chemistry (see our recommended protocol).



Structure of alkyne amidite (aminocyclohexanol, ACH)

General properties

Appearance:	colorless solid
Molecular weight:	409.50
CAS number:	1417539-32-2
Molecular formula:	$C_{21}H_{36}N_3O_3P$
IUPAC name:	trans-4-(5-Hexynoylamino)cyclohexyloxy-N,N-diisopropylamino-2-cyanoethyloxyphosphine
Solubility:	good in acetonitrile and dichloromethane
Quality control:	NMR ^{1}H (95%) and ^{31}P, HPLC-MS
Storage conditions:	Storage: 12 months after receival at -20°C. Transportation: at room temperature for up to 3 weeks. Desiccate.
Legal statement:	This Product is offered and sold for research purposes only. It has not been tested for safety and efficacy in food, drug, medical device, cosmetic, commercial or any other use. Supply does not express or imply authorization to use for any other purpose, including, without limitation, in vitro diagnostic purposes, in the manufacture of food or pharmaceutical products, in medical devices or in cosmetic products.

Oligo synthesis details

Diluent:	acetonitrile
Coupling conditions:	Standard coupling, identical to normal nucleobases.
Cleavage conditions:	standard deprotection
Deprotection conditions:	No deprotection required. Compatible with standard deprotection reagents.