

Lumiprobe Corporation

115 Airport Dr Suite 160 Westminster, Maryland 21157

USA

Phone: +1 888 973 6353 Fax: +1 888 973 6354 Email: order@lumiprobe.com

BDPep 630/650 NHS ester

http://www.lumiprobe.com/p/bdp-630-650-spps-nhs-ester

BDPep 630/650 NHS ester (Dicyano BDP 630/650 NHS ester) is a versatile fluorescent dye with a range of applications in biological and chemical research. The dye is ideal for high-resolution imaging of cellular structures and dynamic processes, providing bright and distinct signals that enhance visualization. Its strong fluorescence allows for effective cell sorting and phenotyping, making it useful in various immunological and cellular studies.


Its remarkable stability in strong acidic TFA media makes it particularly suitable for solid-phase peptide synthesis (SPPS).

Dicyano BDP 630/650 is employed in Förster Resonance Energy Transfer (FRET) assays to investigate molecular interactions and dynamics with high sensitivity. Due to its long excited-state lifetime, this dye is beneficial for detecting binding interactions between molecules, making it valuable in drug discovery and biochemical assays.

The NHS ester form enables efficient labeling of proteins and other biomolecules, that contain a functional amino group. Presence of the C6 spacer allows for more flexibility in conjugation reactions with various biomolecules, making it easier to label proteins and peptides without compromising the dye's fluorescent properties, particularly C6 spacer reduces negative quenching effects.

The dye is typically dissolved in high-quality anhydrous organic solvents such as dimethylformamide (DMF) and dimethylsulfoxide (DMSO), which facilitate its use in conjugation reactions and other applications. Dicyano BDP 630/650 exhibits hydrophobic characteristics, making it less suitable for direct use in aqueous environments compared to more water-soluble dyes like <u>AF 647 NHS ester</u>. Once conjugated to biomolecules, the resulting dicyano BDP 630/650 conjugates can be used in aqueous applications such as fluorescence microscopy and flow cytometry, where they provide reliable fluorescent signals.

BDP 630/650 is characterized by excellent photostability, allowing for prolonged imaging sessions without substantial loss of signal, making it suitable for applications that require extended observation periods.



Structure of BDPep 630/650 NHS ester

Excitation and emission spectra of BDPep 630/650

General properties

Appearance: dark purple powder

Molecular weight: 674.53 Molecular formula: $C_{35}H_{31}BN_6O_6S$

Solubility: good in DMF, DMSO, dichlromethane

Quality control: NMR ¹H and HPLC-MS (95+%)

Storage conditions: 12 months after receival at -20°C in the dark. Transportation: at room temperature

for up to 3 weeks. Desiccate.

Legal statement: Product is offered and sold for research purposes only. Product is not tested for safety

and efficacy in food, drug, medical device, cosmetic, no express or implied authorization to use for any other purpose, including, without limitation, in vitro diagnostic purposes, for humans or animals or for commercial purposes.

Spectral properties

Excitation/absorption maximum, nm: 637 ϵ , L·mol⁻¹·cm⁻¹: 84000 Emission maximum, nm: 663 Fluorescence quantum yield: 0.44 0.17 CF₂₆₀:

CF₂₈₀: 0.18