


## BTAA ligand

<http://www.lumiprobe.com/p/btaa-ligand>

BTAA is the newest generation of water-soluble ligands for Cu(I)-catalyzed azide-alkyne click chemistry reactions (CuAAC). It maintains the Cu(I) oxidation state in the catalyst and protects biomolecules from oxidative damage during the labeling.

BTAA provides much more significant rate enhancement than previous generation ligands (e.g., [THPTA](#) or [TBTA](#)) and suppresses cell cytotoxicity by further lowering copper loading in the catalyst formulation.

A stock solution can be prepared in ddH<sub>2</sub>O and stored at -20°C. Avoid freeze-thaw cycles.



**Structure of BTAA ligand**

### General properties

Appearance:	off-white to grey solid
Molecular weight:	430.52
CAS number:	1334179-85-9
Molecular formula:	C <sub>19</sub> H <sub>30</sub> N <sub>10</sub> O <sub>2</sub>
IUPAC name:	2-(4-((bis((1-(tert-butyl)-1H-1,2,3-triazol-4-yl)methyl)amino)methyl)-1H-1,2,3-triazol-1-yl)acetic acid
Solubility:	Soluble in water, DMSO, DMF, MeOH
Quality control:	NMR <sup>1</sup> H and HPLC-MS (95+%)
Storage conditions:	24 months after receipt at -20°C in the dark. Transportation: at room temperature for up to 3 weeks. Desiccate.