

Lumiprobe Corporation

115 Airport Dr Suite 160 Westminster, Maryland 21157

USA


Phone: +1 888 973 6353 Fax: +1 888 973 6354 Email: order@lumiprobe.com

H-Met-AMC, calpain substrate


http://www.lumiprobe.com/p/h-met-amc-calpain-substrate

H-Met-AMC (L-methionine-7-amido-4-methylcoumarin) is a fluorogenic substrate for methionine aminopeptidase 2 (MetAP2) (Km=310 μ M) and aminopeptidase N (Km=377 μ M). This substrate is a useful tool for inhibitor screening and kinetic analysis.

H-Met-AMC has an excitation peak at 341 nm and emission at 441 nm. The following excitation/emission can also be used: 355,375/440,450.



Structure of H-Met-AMC, calpain substrate



Absorption and emission spectra of calpain substrate

General properties

Appearance: off-white crystals

Molecular weight: 306.39

CAS number: 94367-34-7, 94367-35-8 (trifluoroacetate)

Molecular formula: $C_{15}H_{18}N_2O_3S$

Solubility: Soluble in DMF, DMSO, acetonitrile, ethyl acetate. Slightly soluble in chloroform.

Insoluble in water, diethyl ether

Quality control: NMR ¹H and HPLC-MS (95+%)

Storage conditions: 24 months after receival at -20°C in the dark. Transportation: at room temperature

for up to 3 weeks. Desiccate.

Legal statement: This Product is offered and sold for research purposes only. It has not been tested for

safety and efficacy in food, drug, medical device, cosmetic, commercial or any other use. Supply does not express or imply authorization to use for any other purpose, including, without limitation, in vitro diagnostic purposes, in the manufacture of food

or pharmaceutical products, in medical devices or in cosmetic products.

Spectral properties

Excitation/absorption maximum, nm: 341 Emission maximum, nm: 441